



## ANTIMICROBIAL ACTIVITY TEST OF MEDICINAL PLANT EXTRACT USING ANTIMICROBIAL DISC AND FILTER PAPER AGAINST *Staphylococcus aureus* : CLEAR ZONE RESULTS COMPARISON

Muhammad Taufiq Hidayat<sup>1,\*</sup>, Maharani Pertiwi Koentjoro<sup>2</sup>, Endry Nugroho Prasetyo<sup>3</sup>

<sup>1</sup> Department of Medical Laboratory Technology, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

<sup>2</sup> Department of Economics Development, Faculty of Economics and Business, Universitas Brawijaya, Malang, Indonesia

<sup>3</sup> Department of Biology, Faculty of Science and Data Analytics, Institut Teknologi Sepuluh Nopember, Surabaya, Indonesia

\* Corresponding author : [muhammadtaufiqhidayat7@unusa.ac.id](mailto:muhammadtaufiqhidayat7@unusa.ac.id)

Received : May, 2024

Revised : July, 2024

Accepted : August, 2024

First Publish Online :

August, 20, 2024

Keywords : Antibiotic Test, Antimicrobial Disc, Kirby-Bauer Disc Method, Mannitol Salt Agar, *Staphylococcus aureus*

### ABSTRACT

Plant extracts contain secondary metabolites such as flavonoids and phenolic compounds with antibiotic activity. Antibiotic activity test can be done by using the Kirby-Bauer disc method which the antibiotic activity is indicated by the formation of a clear zone. This research was conducted using extracts of *Moringa oleifera* leaves and *Opuntia cochenillifera* cladode extracted using ethyl acetate as a solvent. Antibiotic activity test against *Staphylococcus aureus* bacteria was done in Mannitol Salt Agar and Luria Bertani Agar media. The Kirby-Bauer discs used are Antimicrobial Disc Oxoid® and Filter Paper. The results of this study show that the use of the Antimicrobial disc Oxoid® and Filter Paper shows no statistical difference in clear zone results (t-value = 0.45; p-value = 0.655; with  $\alpha = 95\%$ ). While the Clear Zone results on Mannitol Salt Agar gave better results than Luria Bertani Agar and were statistically significant (t-value = 2.46; p-value = 0.02; with  $\alpha = 95\%$ ). These findings indicate that filter paper can be an inexpensive alternative for the antibiotic test with no significantly different result compared to commercial antibiotic disc. However, an antibiotic test against *Staphylococcus aureus* is better done on Mannitol Salt Agar compared to Luria Bertani Agar.

This is an open-access article under the [CC-BY-SA](https://creativecommons.org/licenses/by-sa/4.0/) license



## Introduction

*Staphylococcus aureus* is a commensal Gram-positive bacteria that colonize the human nasal mucosa and causes severe infection. The main problem in combating *Staphylococcus aureus* infection is the prevalence of multi-drug resistance, especially due to misuse (excess or abuse)

of antibiotics (Eswari & Yadav, 2019). The increase in antibiotic resistance is estimated to cause the death of up to 10 million cases per year by 2050 (Sharmeen & Fawzi, 2020). Exploration of alternative antibiotics needs to be done to solve the problem of the increase in antibiotic-resistant bacteria. Medicinal plant extract is one promising alternative antibiotics source (Langeveld et

al., 2014). Medicinal plants such as *Moringa oleifera* are widely used for traditional medicine such as to treat dental caries, syphilis, typhoid, diarrhea, epilepsy, purgative, prostate cancer, fever, and HIV-AIDS (Dzotam et al., 2016). *Moringa oleifera* contains secondary metabolites including alkaloids, flavonoids (quercetin and kaempferol) (Lin et al., 2018; Wang et al., 2017), saponins, and tannins (Sulastri et al., 2018) which have antibiotic activity (Ilanko et al., 2019). Other used medicinal plant in this study, *Opuntia cochenillifera*, contain phenolic compounds, flavonoids, and anthocyanins (Alves et al., 2017) with antibacterial activity. Cladode is part of *Opuntia cochenillifera* that contain alkaloids, saponin, tannins, terpenes, flavonoid, and hydroxi flavones (Monrroy et al., 2017). One of the mechanisms of action of antibiotics from plant extract is through inhibition of peptidoglycan synthesis. Many of the proteins involved in this pathway are Mur enzymes and the penicillin-binding proteins (PBPs), which are known as good targets for antibiotics (Liu & Breukink, 2016; Tomoda, 2016).

One of the important things to determine the antibiotic activity of a compound is the method of determining antibiotic activity. Kirby-Bauer disc method is one of the major methods to test antibiotic susceptibility. The Kirby-Bauer disc method relies on the diffusion of the test substance from the filter discs to the bacterial cultures (Kourmouli et al., 2018). The disk diffusion method is included in the agar diffusion method because the plant extract diffuses from its reservoir through the agar medium. The reservoir (filter paper disc) is placed on top of an agar surface. If tested plant extracts compounds have antibiotic activity, the inhibition zone will be developed around the filter paper disk after incubation. The diameter of the inhibition zone describes the antimicrobial potency of plant extracts (Horváth et al., 2016). One of the most commonly used discs for antibiotic susceptibility test is the

commercial Antimicrobial Disc. In this study, we tested the use of inexpensive filter paper as an alternative to commercial antimicrobial discs.

Another factors that influence the clear zone results are the media used when testing antibiotic activity. The agar media must allow free diffusion of the antimicrobial from the disc. Mannitol Salt Agar (MSA) is applied to isolate and identify *Staphylococcus aureus* as a differential and selective medium. Mannitol Salt Agar contain peptones and beef extract, both of which are necessary for growth. The 7.5% sodium chloride concentration inhibits bacterial organisms other than *staphylococci* partially or completely. Mannitol fermentation (in MSA) produces acid, which is detectable by phenol red indicator and assists in the separation of *staphylococcal* species. Coagulase-positive *staphylococci* (e.g., *Staphylococcus aureus*) create yellow colonies and a yellow medium surrounding them (Aryal, 2019). Luria Bertani Agar consists of tryptone, yeast extract, NaCl, and agar (Macwilliams & Liao, 2006). This study aimed to compare the use of commercial Antimicrobial Disc and Filter Paper, as well as the use of Mannitol Salt agar and Luria Bertani Agar media in susceptibility testing of medicinal plant extracts against *Staphylococcus aureus*.

## Materials and Methods

### Plant Extraction

The plant extraction process is done based on (Kher et al., 2019) with modifications. *Moringa oleifera* leaves and *Opuntia cochenillifera* cladode was extracted using ethyl acetate as a solvent. The plant samples were oven-dried at 50 °C until they reached a constant weight. After drying, the sample is crushed into a powder. Then, the plant samples were macerated using ethyl acetate solvent with a mass ratio of plant samples to the solvent

volume of 1:10 for 24 hours. In a rotary evaporator, the filtrate is concentrated until crude extract is obtained.

#### Inoculation of *Staphylococcus aureus* to agar media

Inoculation of *Staphylococcus aureus* was done using the swab technique based on (Weme, 2018) with modifications. One loop from a single colony of *S. aureus* culture was inoculated aseptically into Luria Bertani Broth media then incubated for 24 hours at 37° C. The sterile cotton is dipped in overnight-culture broth, then swab evenly over the surface of the agar medium.

#### Kirby-Bauer Antimicrobial Susceptibility Test

The antimicrobial activity assay is done based on (Yu et al., 2019) with modifications. Each of the Antimicrobial Disc Oxoid® and Filter Paper with a diameter of 5 mm was immersed in a plant extract with a concentration of 50 mg/mL, 25 mg/mL, 12.5 mg/mL, and 6.25 mg/mL, and amoxicillin 25 mg/mL as a positive control and DMSO 1% as a negative control. After immersing for 1 minute, each disc was placed on the surface of the media that had been inoculated with *S. aureus*. Culture on solid media with a disc was incubated for 24 hours at 37°C.

#### Statistical Analysis

Each plate consists of 6 test points that have been given a number code (1-6)

consisting of plant extracts of 50 mg/mL (1), 25 mg/mL (2), 12.5 mg/mL (3), 6.25 mg/mL (4), amoxicillin 25 mg/mL (5), and DMSO 1% (6). The data entered are test points number 1 to 5, while negative control is not included in the statistical test because as a negative control it should be 0 in clear zone diameter. DMSO is widely used as a solvent for antibiotic compounds (Camp et al., 2020) and is known to have low toxicity (Phaechamud & Setthajindalert, 2018). Each test consists of 2 plates for *Moringa oleifera* leaves and 1 plate for *Opuntia cochenillifera* cladode so that each treatment consisted of 15 data replications. The t-test statistical test was used to compare whether the clear zone diameter value between the two treatments was significantly different or not. Statistical analysis was done using Minitab® with a confidence level of 95%.

## Results and Discussion


### *Comparison of using Commercial Antimicrobial Disc and Filter Paper*

Disc diffusion is an excellent, low-cost, and versatile method for Antibiotic Susceptibility Testing, and its quality is dictated by the discs and media used (Åhman et al., 2019). The clear zone results after 24 hours incubation (Table 1) showed that the use of filter paper and commercial Antimicrobial Disc was not statistically significant (T-Value = 0.45 P-Value = 0.655 with  $\alpha = 95\%$ )

Table 1 Clear Zone Comparison of Disc

	Filter Paper	Antimicrobial Disc Oxoid®
Clear Zone Average (mm)	11,27±4,91	12,07±4,79

Image



Statistical  
Analysis

T-Value = 0,45 P-Value = 0,655 with  $\alpha = 95\%$

The commercial Oxoid Antibiotic discs are made from paper which conforms to the WHO and FDA standards (Joshi et al., 2008). The plant extract diffuses from the disc into the agar media inoculated with the test microorganism during the incubation phase. The active antimicrobial extracts or chemicals cause zones of inhibition to form around the disc, providing the value of the minimum inhibitory concentration (MIC). Bioactive chemicals found in *Moringa oleifera* include 9- Octadecenoic acid (z) -, Heptadecanoic acid, and phytol acetate (Syeda & Riazunnisa, 2020). *Opuntia cochenillifera* cladode contains flavonoids, phenols, alkaloids, tannins, and saponins (Suryawanshi & Vidyasagar, 2016).

The diameter of the filter paper disk, the amount of substance placed on the disk, the kind and concentration of agar, the medium depth and pH, the bacterial strain tested, and the incubation temperature are all parameters that determine the size of the inhibitory zones (Horváth et al., 2016). The average clear zone value of the filter paper ( $11.27 \pm 4.91$  mm) was smaller than the Antimicrobial Disc Oxoid® ( $12.07 \pm 4.79$  mm) but did not differ significantly.



Antimicrobial Disc Oxoid® is made of unique inert material which enhances their absorption hence allowing faster adhesion of discs to the media (Justesen et al., 2013). The active compounds of the extract are less retained in the filter paper than a commercial antimicrobial disc (Royo et al., 2010) so that in this study, 2 sheets of filter paper were placed on the media. Research from (Ogba et al., 2018) shows that locally prepared discs using Whatman filter paper 3 are more effective than commercial discs. These results indicate that the use of filter paper (2 layers) can provide clear zone results that are not significantly different from commercial Antimicrobial Discs. Filter paper can be an inexpensive alternative to commercial Antimicrobial Discs with results that are not significantly different.

#### *Comparison of using Mannitol Salt Agar and Luria Bertani Agar*

Media is one of the factors that influence clear zone results (Åhman et al., 2019). The results of this study indicate that the use of Mannitol Salt Agar is better than Luria Bertani Agar for Antibiotic Susceptibility for *Staphylococcus aureus*.



Table 2. Clear Zone Comparison of Media

	Mannitol Salt Agar	Luria Bertani Agar
Clear Zone Average (mm)	13,47±3,81	9,73±4,46
Image		
Statistical Analysis	T-Value = 2,46 P-Value = 0,020 with $\alpha = 95\%$	

Clear Zone results on Mannitol Salt Agar gave better results than Luria Bertani Agar and were statistically significant (t-value = 2.46; p-value = 0.02; with  $\alpha = 95\%$ ). For many years, MSA has been employed as a selective, differential media for the isolation of *S. aureus* (Sharp & Searcy, 2006). Several factors that affect the accuracy and repeatability of disc diffusion methods include bacterial inoculum preparation, manual streaking of media plates, disk content, agar medium, nutritional requirements, incubation temperature and atmosphere, incubation time, and subjectivity of inhibition zone reading (Cherkaoui et al., 2020). Mannitol fermentation (in MSA) produces acid, which is detectable by phenol red indicator and facilitates in the distinguishing of *staphylococcal* species. Coagulase-positive staphylococci (e.g., *Staphylococcus aureus*) generate yellow colonies and a yellow medium surrounding them (Aryal, 2019). The 7.5% sodium chloride concentration inhibits bacterial organisms other than staphylococci partially or completely. Luria Bertani Agar consists of tryptone, yeast extract, NaCl, and agar (Macwilliams & Liao, 2006). The use of MSA will limit the contamination of other bacteria than

*Staphylococcus aureus*. The results of this study indicate that the use of selective media provides the best clear zone results for the Antibiotic Susceptibility test. Mannitol salt agar is the right medium for the Antibiotic Susceptibility test for *Staphylococcus aureus*.

### Conclusions

In conclusions, we find that the use of the Antimicrobial disc Oxoid® and Filter Paper shows no statistical difference in clear zone results (t-value = 0.45; p-value = 0.655; with  $\alpha = 95\%$ ). While the Clear Zone results on Mannitol Salt Agar gave better results than Luria Bertani Agar and were statistically significant (t-value = 2.46; p-value = 0.02; with  $\alpha = 95\%$ ). These findings indicate that filter paper can be an inexpensive alternative for the antibiotic test with no significantly different result compared to commercial antibiotic disc. However, an antibiotic test against *Staphylococcus aureus* is better done on Mannitol Salt Agar compared to Luria Bertani Agar.

## Acknowledgment

This study was supported by the Ministry of Research, and Technology / National Research and Innovation Agency (RISTEKBRIN), Republic of Indonesia, and Institut Teknologi Sepuluh Nopember (ITS), Surabaya, Indonesia under a grant number of 3/AMD/E1/KP.PTNBH/2020 and 1177/PKS/ITS/2019.

## References

- Åhman, J., Matuschek, E., & Kahlmeter, G. (2019). The quality of antimicrobial discs from nine manufacturers—EUCAST evaluations in 2014 and 2017. *Clinical Microbiology and Infection*, 25(3), 346–352. <https://doi.org/10.1016/j.cmi.2018.05.021>
- Alves, F. A. L., De Andrade, A. P., Bruno, R. de L. A., Silva, M. G. de V., De Souza, M. de F. V., & Dos Santos, D. C. (2017). Seasonal variability of phenolic compounds and antioxidant activity in prickly pear cladodes of *Opuntia* and *Nopalea* genres. *Food Science and Technology*, 37(4), 536–543. <https://doi.org/10.1590/1678-457x.19316>
- Aryal, S. (2019). *Mannitol Salt Agar for the isolation of Staphylococcus aureus*. <https://microbiologyinfo.com/mannitol-salt-agar-for-the-isolation-of-staphylococcus-aureus/>
- Camp, J. E., Nyamini, S. B., & Scott, F. J. (2020). Cyrene™ is a green alternative to DMSO as a solvent for antibacterial drug discovery against ESKAPE pathogens. *RSC Medicinal Chemistry*, 11(1), 111–117. <https://doi.org/10.1039/c9md00341j>
- Cherkaoui, A., Renzi, G., Fischer, A., Azam, N., Schorderet, D., Vuilleumier, N., & Schrenzel, J. (2020). Comparison of the Copan WASPLab incorporating the BioRad expert system against the SIRscan 2000 automatic for routine antimicrobial disc diffusion susceptibility testing. *Clinical Microbiology and Infection*, 26(5), 619–625. <https://doi.org/10.1016/j.cmi.2019.11.008>
- Dzotam, J. K., Touani, F. K., & Kuete, V. (2016). Antibacterial and antibiotic-modifying activities of three food plants (*Xanthosoma mafaffa* Lam., *Moringa oleifera* (L.) Schott and *Passiflora edulis* Sims) against multidrug-resistant (MDR) Gram-negative bacteria. *BMC Complementary and Alternative Medicine*, 16(1), 1–8. <https://doi.org/10.1186/s12906-016-0990-7>
- Eswari, J. S., & Yadav, M. (2019). South African Journal of Botany New perspective of drug discovery from herbal medicinal plants : *Andrographis paniculata* and *Bacopa monnieri* ( terpenoids ) and novel target identification against *Staphylococcus aureus*. *South African Journal of Botany*, 124, 188–198. <https://doi.org/10.1016/j.sajb.2019.05.013>
- Horváth, G., Bencsik, T., Ács, K., & Kocsis, B. (2016). Sensitivity of ESBL-Producing Gram-Negative Bacteria to Essential Oils, Plant Extracts, and Their Isolated Compounds. In *Antibiotic Resistance: Mechanisms and New Antimicrobial Approaches* (Issue 2). <https://doi.org/10.1016/B978-0-12-803642-6.00012-5>
- Ilanko, P., McDonnell, P. A., van Vuuren, S., & Cock, I. E. (2019). Interactive antibacterial profile of *Moringa oleifera* Lam. extracts and conventional antibiotics against bacterial triggers of some autoimmune inflammatory diseases. *South African Journal of Botany*, 124, 420–435. <https://doi.org/10.1016/j.sajb.2019.04.008>

- Joshi, A., Iyer, V., Balasubramaniam, U., Kagal, A., & Bharadwaj, R. (2008). Comparison of efficacy of three commercially available antibiotic discs. *Indian Journal of Medical Microbiology*, 26(2), 160–162. <https://doi.org/10.4103/0255-0857.40533>
- Justesen, U. S., Acar, Z., Olsson, K., Jensen, T. G., Kern, M. B., Skov, R. L., & Gahrn-Hansen, B. (2013). Comparison of Rosco Neo-Sensitabs with Oxoid paper disks in EUCAST disk diffusion antimicrobial susceptibility testing on Mueller-Hinton agar. *European Journal of Clinical Microbiology and Infectious Diseases*, 32(5), 621–625. <https://doi.org/10.1007/s10096-012-1785-5>
- Kher, M. N., Sheth, N. R., & Bhatt, V. D. (2019). In Vitro Antibacterial Evaluation of Terminalia chebula as an Alternative of Antibiotics against Bovine Subclinical Mastitis. *Animal Biotechnology*, 30(2), 151–158. <https://doi.org/10.1080/10495398.2018.1451752>
- Kourmouli, A., Valenti, M., van Rijn, E., Beaumont, H. J. E., Kalantzi, O. I., Schmidt-Ott, A., & Biskos, G. (2018). Can disc diffusion susceptibility tests assess the antimicrobial activity of engineered nanoparticles? *Journal of Nanoparticle Research*, 20(3), 2–7. <https://doi.org/10.1007/s11051-018-4152-3>
- Langeveld, W. T., Veldhuizen, E. J. A., & Burt, S. A. (2014). Synergy between essential oil components and antibiotics: A review. *Critical Reviews in Microbiology*, 40(1), 76–94. <https://doi.org/10.3109/1040841X.2013.763219>
- Lin, M., Zhang, J., & Chen, X. (2018). Bioactive flavonoids in Moringa oleifera and their health-promoting properties. *Journal of Functional Foods*, 47(June), 469–479. <https://doi.org/10.1016/j.jff.2018.06.011>
- Liu, Y., & Breukink, E. (2016). The Membrane Steps of Bacterial Cell Wall Synthesis as Antibiotic Targets. *Antibiotics*, 5(28), 1–22. <https://doi.org/10.3390/antibiotics5030028>
- Macwilliams, M. P., & Liao, M. (2006). Luria Broth (LB) and Luria Agar (LA) Media and Their Uses Protocol Resource Type: Publication Date: Authors. *American Society for Microbiology*, October 2006, 7–9.
- Monrroy, M., García, E., Ríos, K., & García, J. R. (2017). Extraction and Physicochemical Characterization of Mucilage from Opuntia cochenillifera (L.) Miller. *Journal of Chemistry*, 2017, 1–9. <https://doi.org/10.1155/2017/4301901>
- Ogba, O. M., Udo, N. I., Inyang-etoh, P. C., & Olorode, O. A. (2018). Evaluation of the Potency of Commercial and Locally Prepared Evaluation of the Potency of Commercial and Locally Prepared Antibiotic Discs on Clinical Bacterial Isolates in Calabar, Nigeria. *International Journal of Biomedicine & Life Sciences*, 2(1), 1–7. <https://doi.org/10.5281/zenodo.1119113> EVALUATION
- Phaechamud, T., & Setthajindalert, O. (2018). Antimicrobial in-situ forming gels based on bleached shellac and different solvents. *Journal of Drug Delivery Science and Technology*, 46(March), 285–293. <https://doi.org/10.1016/j.jddst.2018.05.035>
- Royo, M., Fernández-Pan, I., & Maté, J. I. (2010). Antimicrobial effectiveness of oregano and sage essential oils incorporated into whey protein films or cellulose-based filter paper. *Journal of the Science of Food and Agriculture*, 90(9), 1513–1519. <https://doi.org/10.1002/jsfa.3977>

- Sharmeen, B., & Fawzi, M. (2020). Essential oils from 9 exotic and endemic medicinal plants from Mauritius shows in vitro antibacterial and antibiotic potentiating activities. *South African Journal of Botany*, *132*, 355–362.  
<https://doi.org/10.1016/j.sajb.2020.05.001>
- Sharp, S. E., & Searcy, C. (2006). Comparison of mannitol salt agar and blood agar plates for identification and susceptibility testing of *Staphylococcus aureus* in specimens from cystic fibrosis patients. *Journal of Clinical Microbiology*, *44*(12), 4545–4546.  
<https://doi.org/10.1128/JCM.01129-06>
- Sulastri, E., Zubair, M. S., Anas, N. I., Abidin, S., Hardani, R., Yulianti, R., Sulastri, E., Zubair, S., Anas, I., & Hardani, R. (2018). Total Phenolic , Total Flavonoid , Quercetin Content and Antioxidant Activity of Standardized Extract of *Moringa oleifera* Leaf from Regions with Different Elevation. *Pharmacognosy Journal*, *10*(6), 104–108.  
<https://doi.org/10.5530/pj.2018.6s.20>
- Suryawanshi, P. K., & Vidyasagar, G. M. (2016). Phytochemical Screening for Secondary Metabolites of *Opuntia cochenillifera* ( L.) Mill. *International Journal of Life Sciences Research*, *4*(2), 145–151.
- Syeda, A. M., & Riazunnisa, K. (2020). Data on GC-MS analysis, in vitro anti-oxidant and anti-microbial activity of the *Catharanthus roseus* and *Moringa oleifera* leaf extracts. *Data in Brief*, *29*, 105258.  
<https://doi.org/10.1016/j.dib.2020.105258>
- Tomoda, H. (2016). Recent Progress in Study on the Biologically-Active Natural Products New Approaches to Drug Discovery for Combating MRSA. *Chem. Pharm. Bull.*, *64*(2), 104–111.  
<https://doi.org/10.1248/cpb.c15-00743>
- Wang, Y., Gao, Y., Ding, H., Liu, S., Han, X., Gui, J., & Liu, D. (2017). Subcritical ethanol extraction of flavonoids from *Moringa oleifera* leaf and evaluation of antioxidant activity. *Food Chemistry*, *218*, 152–158.  
<https://doi.org/10.1016/j.foodchem.2016.09.058>
- Weme, E. T. (2018). Rapid antimicrobial susceptibility testing of positive blood cultures by direct inoculation and reading of disc diffusion tests after 3–4 hours. *Apmis*, *126*(11), 870–876.  
<https://doi.org/10.1111/apm.12897>
- Yu, H., Liu, M., Liu, Y., Qin, L., Jin, M., & Wang, Z. (2019). Antimicrobial activity and mechanism of action of *Dracocephalum moldavica* L. extracts against clinical isolates of *Staphylococcus aureus*. In *Frontiers in Microbiology* (Vol. 10, Issue JUN).  
<https://doi.org/10.3389/fmicb.2019.01249>